

Pharmacological Considerations for Pregnant Women

The pharmacological agents listed below are to be used only for indicated medical conditions and with appropriate supervision.

Pharmaceutical Agent	Indications, Contraindications, and Special Considerations
Analgesics	
Acetaminophen	May be used during pregnancy. Oral pain can often be managed with non-opioid medication. If opioids are used, prescribe the lowest dose for the shortest duration (usually less than 3 days), and avoid issuing refills to reduce risk for dependency.
Acetaminophen with Codeine, Hydrocodone, or Oxycodone	
Codeine	
Meperidine	
Morphine	
Aspirin	May be used in short duration during pregnancy; 48 to 72 hours. Avoid in 1st and 3rd trimesters.
Ibuprofen	
Naproxen	
Antibiotics	
Amoxicillin	May be used during pregnancy.
Cephalosporins	
Clindamycin	
Metronidazole	
Penicillin	
Ciprofloxacin	Avoid during pregnancy.
Clarithromycin	
Levofloxacin	
Moxifloxacin	
Tetracycline	Never use during pregnancy.
Anesthetics	
	Consult with a prenatal care health professional before using intravenous sedation or general anesthesia. Limit duration of exposure to less than 3 hours in pregnant women in the third trimester.
Local anesthetics with epinephrine (e.g., Bupivacaine, Lidocaine, Mepivacaine)	May be used during pregnancy.
Nitrous oxide (30%)	May be used during pregnancy when topical or local anesthetics are inadequate. Pregnant women require lower levels of nitrous oxide to achieve sedation; consult with prenatal care health professional.
Antimicrobials	
	Use alcohol-free products during pregnancy.
Cetylpyridinium chloride mouth rinse	May be used during pregnancy.
Chlorhexidine mouth rinse	
Xylitol	

From *Oral Health Care During Pregnancy: A National Consensus Statement—Summary of an Expert Workgroup Meeting* © 2012 by the National Maternal and Child Oral Health Resource Center, Georgetown University. Table updated 2017. Permission is given to photocopy this publication or to forward it, in its entirety, to others.